

# Two-Weighted Inequalities for Maximal Commutators in Generalized Weighted Morrey Spaces on Spaces of Homogeneous Type

Ayşenur Aydoğdu<sup>1</sup>, Canay Aykol<sup>1,†</sup> and Javanshir J. Hasanov<sup>2</sup>

1. Department of Mathematics, Ankara University, Ankara, Turkey

2. Azerbaijan State Oil and Industry University, Baku, Azerbaijan

<sup>†</sup>Corresponding author

## Abstract

In this paper we give a characterization of two-weighted inequalities for maximal commutators in generalized weighted Morrey spaces on spaces of homogeneous type  $\mathcal{M}_\omega^{p,\varphi}(X)$ . We prove the boundedness of maximal commutators  $[M, b]$  from the spaces  $\mathcal{M}_{\omega_1}^{p,\varphi_1}(X)$  to the spaces  $\mathcal{M}_{\omega_2}^{p,\varphi_2}(X)$ , where  $1 < p < \infty$ ,  $0 < \delta < 1$  and  $(\omega_1, \omega_2) \in \tilde{A}_p(X)$ .

**Keywords:** Maximal operator; commutator; generalized weighted Morrey space; spaces of homogeneous type; regularity

**2020 Mathematics Subject Classification:** 42B20, 42B25, 42B35

**Article history:** Received 13 Jun 2023; Revised 3 Aug 2023; Accepted 10 Aug 2023; Online 9 Sep 2023

R. Coifman and G. Weiss introduced certain topological measure spaces which are equipped with a metric which is compatible with the given measure in a sense in the 1970s. These spaces are called spaces of homogeneous type. In this work, we find necessary and sufficient conditions for the boundedness of Hardy-Littlewood maximal commutators in generalized weighted Morrey spaces on spaces of homogeneous type. As a generalization of  $L_p(\mathbb{R}^n)$ , the classical Morrey spaces were introduced by Morrey [30] to study the local behavior of solutions to second-order elliptic partial differential equations. Moreover, various Morrey spaces are defined in the process of study. Guliyev, Mizuhara and Nakai [15, 29, 32] introduced generalized Morrey spaces  $M^{p,\varphi}(\mathbb{R}^n)$  (see, also [16, 17, 38]).

Recently, Komori and Shirai [27] defined the weighted Morrey spaces  $L_w^{p,\kappa}(\mathbb{R}^n)$  and studied the boundedness of some classical operators such as the Hardy-Littlewood maximal operator, the Calderón-Zygmund operator on these spaces. Also, Guliyev in [18] first introduced the generalized weighted Morrey spaces  $M_w^{p,\varphi}(\mathbb{R}^n)$  and studied the boundedness of the sublinear

**Contact:** Ayşenur Aydoğdu ✉ [anaydogdu@ankara.edu.tr](mailto:anaydogdu@ankara.edu.tr); Canay Aykol ✉ [canay.Aykol@science.ankara.edu.tr](mailto:canay.Aykol@science.ankara.edu.tr); Javanshir J. Hasanov ✉ [hasanovjavanshir@gmail.com](mailto:hasanovjavanshir@gmail.com).

© 2023 The Author(s). Published by Mersin University Press. This is an Open Access article distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

operators and their higher order commutators generated by Calderón-Zygmund operators and Riesz potentials in these spaces (see, also [21, 25]). Note that, Guliyev [18] gave the concept of generalized weighted Morrey space which could be viewed as an extension of both  $M_w^{p,\varphi}(\mathbb{R}^n)$  and  $L_w^{p,\kappa}(\mathbb{R}^n)$ .

We say that  $X = (X, d, \mu)$  is a space of homogeneous type in the sense of Coifman and Weiss [10] if  $d$  is a quasi-metric on  $X$  and  $\mu$  is a positive measure satisfying the doubling condition, i.e.  $X$  is a topological space endowed with a quasi-metric  $d$  and a positive measure  $\mu$  such that

$$\begin{aligned} d(x, y) &= d(y, x) \geq 0 \text{ for all } x, y \in X, \\ d(x, y) &= 0 \text{ if and only if } x = y, \\ d(x, y) &\leq C_k [d(x, z) + d(z, y)] \text{ for all } x, y, z \in X, \end{aligned}$$

the balls  $B(x, r) = \{y \in X : d(x, y) < r\}$ ,  $r > 0$ , form a basis of neighborhoods of point  $x$ ,  $\mu$  is defined on a  $\sigma$ -algebra of subsets of  $X$  which contains the balls, and

$$0 < \mu(B(x, 2r)) < C_\mu \mu(B(x, r)) < \infty,$$

where  $C_k, C_\mu \geq 1$  are constants independent of  $x, y, z \in X$  and  $r > 0$ . As usual, the dilation of a ball  $B = B(x, r)$  will be denoted by  $\lambda B = B(x, \lambda r)$  for every  $\lambda > 0$ .

Throughout this paper we always assume that  $\mu(X) = \infty$ , the space of compactly supported continuous function is dense in  $L_1(X, \mu)$  and that  $X$  is  $N$ -homogeneous ( $N > 0$ ), i.e.

$$C_1 r^N \leq \mu(B(x, r)) \leq C_2 r^N,$$

where  $C_i \geq 1$  ( $i = 1, 2$ ) are constants independent of  $x$  and  $r$ . The  $n$ -dimensional Euclidean space  $\mathbb{R}^n$  is  $n$ -homogeneous.

Let  $(X, d, \mu)$  be a homogeneous space,  $1 \leq p < \infty$ ,  $\varphi$  be a positive measurable function on  $(0, \infty)$  and  $\omega$  be a non-negative measurable function on  $X$ . We denote by  $\mathcal{M}_\omega^{p,\varphi}$  the generalized weighted Morrey space on spaces of homogeneous type, the space of all functions  $f \in L_{p,\omega}^{loc}(X)$  with finite norm

$$\|f\|_{\mathcal{M}_\omega^{p,\varphi}} = \sup_{x \in X, r > 0} \frac{1}{\varphi(x, r) \|\omega\|_{L_p(B(x, r))}} \|f\|_{L_{p,\omega}(B(x, r))},$$

where the supremum is taken over all balls  $B(x, r)$  in  $X$  and  $L_{p,\omega}(B(x, r))$  denotes the weighted  $L_p$ -space of measurable functions  $f$  for which

$$\|f\|_{L_{p,\omega}(B(x, r))} \equiv \|f \chi_{B(x, r)}\|_{L_{p,\omega}(X)} = \left( \int_{B(x, r)} |f(y)|^p \omega(y) d\mu(y) \right)^{\frac{1}{p}}.$$

Moreover, by  $W\mathcal{M}_\omega^{p,\varphi}$  we denote the weak generalized weighted Morrey space on spaces of homogeneous type of all functions  $f \in WL_{p,\omega}^{loc}(X)$  with finite norm

$$\|f\|_{W\mathcal{M}_\omega^{p,\varphi}} = \sup_{x \in X, r > 0} \frac{1}{\varphi(x, r) \|\omega\|_{L_p(B(x, r))}} \|f\|_{WL_{p,\omega}(B(x, r))},$$

where  $WL_{p,\omega}(B(x, r))$  denotes the weak weighted  $L_p$ -space of measurable functions  $f$  for which

$$\|f\|_{WL_{p,\omega}(B(x, r))} \equiv \|f \chi_{B(x, r)}\|_{WL_{p,\omega}(X)} = \sup_{t > 0} t \left( \int_{\{y \in B(x, r) : |f(y)| > t\}} |f(y)|^p \omega(y) d\mu(y) \right)^{\frac{1}{p}}.$$

Note that if  $\omega(x) = \chi_{B(x,r)}$ , then  $\mathcal{M}_\omega^{p,q}(X) = \mathcal{M}^{p,q}(X)$  is the generalized Morrey space and if  $\varphi(x,r) = \left(\frac{r^\lambda}{\mu(B(x,r))}\right)^{\frac{1}{p}}$ , then  $\mathcal{M}_\omega^{p,q}(X) = L_{p,\lambda}(X)$  is the classic Morrey space.

We now recall the definition of Hardy-Littlewood operator on space of homogeneous type.

Let  $f$  be a locally integrable function on  $X$ . The so-called of Hardy-Littlewood maximal function is defined by the formula

$$Mf(x) = \sup_{r>0} \frac{1}{\mu(B(x,r))} \int_{B(x,r)} |f(y)| d\mu(y),$$

where  $\mu(B(x,r))$  is measure of the ball  $B(x,r)$ .

The study of maximal operators is one of the most important topics in harmonic analysis. These significant non-linear operators, whose behavior are very informative in particular in differentiation theory, provided the understanding and the inspiration for the development of the general class of singular and potential operators.

In this paper we aim to give a characterization of two-weighted inequalities for maximal commutators in generalized weighted Morrey spaces on spaces of homogeneous type. Two-weight norm inequalities for fractional maximal operators and singular integrals on Lebesgue spaces were widely studied (see, for example [11, 12, 14, 26, 28]). The weighted norm inequalities with different types of weights on Morrey spaces were also studied (see, for example [22, 33, 37]). The two-weight norm inequalities for maximal operator, fractional maximal operator, singular integral operators, Riesz potential and fractional integral operators on Morrey-type spaces were studied by many authors (see, for example [3, 7, 8, 35, 39]). Also, two-weighted inequalities for maximal operator in generalized weighted Morrey spaces on spaces of homogeneous type were studied in [4].

In the sequel we use the letter  $C$  for a positive constant, independent of appropriate parameters and not necessary the same at each occurrence. For every  $p \in [1, \infty]$ , we denote  $p'$  the conjugate of  $p$ , i.e.,  $\frac{1}{p} + \frac{1}{p'} = 1$ .  $\mathfrak{M}(\mathbb{R}_+)$ ,  $\mathfrak{M}^+(\mathbb{R}_+)$  and  $\mathfrak{M}^+(\mathbb{R}_+; \uparrow)$  stand for the set of Lebesgue-measurable functions on  $\mathbb{R}_+$ , and its subspaces of nonnegative and nonnegative non-decreasing functions, respectively.

## 1 Preliminaries

Let  $(X, d, \mu)$  be space of  $N$ -homogeneous type as mentioned in Section 1. We now recall the definition of  $A_p$  weight functions.

**Definition 1.1.** The weight function  $\omega$  belongs to the class  $A_p(X)$  for  $1 \leq p < \infty$ , if

$$\sup_{x \in X, r > 0} \left( \mu(B(x,r))^{-1} \int_{B(x,r)} \omega^p(y) d\mu(y) \right)^{\frac{1}{p}} \left( \mu(B(x,r))^{-1} \int_{B(x,r)} \omega^{-p'}(y) d\mu(y) \right)^{\frac{1}{p'}}$$

is finite and  $\omega$  belongs to  $A_1(X)$ , if there exists a positive constant  $C$  such that for any  $x \in X$  and  $r > 0$

$$\mu(B(x,r))^{-1} \int_{B(x,r)} \omega(y) d\mu(y) \leq C \operatorname{ess\,sup}_{y \in B(x,r)} \frac{1}{\omega(y)}.$$

The weight function  $(\omega_1, \omega_2)$  belongs to the class  $\tilde{A}_p(X)$  for  $1 < p < \infty$ , if

$$\sup_{x \in X, r > 0} \left( \mu(B(x, r))^{-1} \int_{B(x, r)} \omega_2^p(y) d\mu(y) \right)^{\frac{1}{p}} \left( \mu(B(x, r))^{-1} \int_{B(x, r)} \omega_1^{-p'}(y) d\mu(y) \right)^{\frac{1}{p'}}$$

is finite.

The following theorem was proved in [31] (Theorem 1, p. 209 and Theorem 2, p. 215).

**Theorem 1.2.** Let  $1 \leq p < \infty$ .

- 1) Then the operator  $M$  is bounded in  $L_{p, \omega}(X)$  if and only if  $\omega \in A_p(X)$ .
- 2) Then the operator  $M$  is bounded from  $L_{1, \omega}(X)$  to  $WL_{1, \omega}(X)$  if and only if  $\omega \in A_1(X)$ .

**Theorem 1.3.** [4] Let  $1 < p < \infty$ ,  $0 < \delta < 1$  and  $(\omega_1, \omega_2) \in \tilde{A}_p(X)$ . Then

$$\|Mf\|_{L_{p, \omega_2^\delta}(B(x, r))} \leq C \|\omega_2^\delta\|_{L_p(B(x, r))} \sup_{t > r} \|f\|_{L_{p, \omega_1^\delta}(B(x, t))} \|\omega_2^\delta\|_{L_p(B(x, t))}^{-1}$$

for every  $f \in L_{p, \omega_1^\delta}(X)$ , where  $C$  does not depend on  $f, x$  and  $r$ .

**Theorem 1.4.** [4] Let  $1 < p < \infty$ ,  $0 < \delta < 1$ ,  $(\omega_1, \omega_2) \in \tilde{A}_p(X)$  and the function  $\varphi_1(x, r)$  and  $\varphi_2(x, r)$  satisfy the condition

$$\sup_{t > r} \frac{\text{ess inf}_{t < s < \infty} \varphi_1(x, s) \|\omega_1^\delta\|_{L_p(B(x, s))}}{\|\omega_2^\delta\|_{L_p(B(x, t))}} \leq C \varphi_2(x, r),$$

where  $C$  does not depend on  $x$  and  $t$ .

Then the operator  $M$  is bounded from the space  $\mathcal{M}_{\omega_1^\delta}^{p, \varphi_1}(X)$  to the space  $\mathcal{M}_{\omega_2^\delta}^{p, \varphi_2}(X)$ .

**Lemma 1.5.** [34] Let  $1 < p < \infty$  and  $(\omega_1, \omega_2) \in \tilde{A}_p(X)$ , then  $(\omega_2^{-1}, \omega_1^{-1}) \in \tilde{A}_{p'}(X)$ , with  $\frac{1}{p} + \frac{1}{p'} = 1$ .

**Lemma 1.6.** [34] Let  $1 < p < \infty$ ,  $0 < \delta < 1$  and  $(\omega_1, \omega_2) \in \tilde{A}_p(X)$ . If  $\frac{q-1}{p-1} = \delta$ , then  $(\omega_1, \omega_2) \in \tilde{A}_{q'}(X)$ , with  $\frac{1}{p} + \frac{1}{p'} = 1$ .

**Corollary 1.7.** [34] Let  $1 < p < \infty$ ,  $0 < \delta < 1$  and  $(\omega_1, \omega_2) \in \tilde{A}_p(X)$ , then the operator  $M$  is bounded from  $L_{p, \omega_1^\delta}(X)$  to  $L_{p, \omega_2^\delta}(X)$ .

Let  $M^\sharp$  be the sharp maximal function defined by

$$M^\sharp f(x) = \sup_{r > 0} \mu(B(x, r))^{-1} \int_{B(x, r)} |f(y) - f_{B(x, r)}| d\mu(y),$$

where  $f_{B(x, r)}(x) = \mu(B(x, r))^{-1} \int_{B(x, r)} f(y) d\mu(y)$ .

**Lemma 1.8.** Let  $1 < p < \infty$  and  $\omega \in A_p(X)$ . Then

$$\|f\omega\|_{L_p} \leq C \|\omega M^\sharp f\|_{L_p}$$

with a constant  $C > 0$  not depending on  $f$ .

**Definition 1.9.** We define the  $BMO(X)$  space as the set of all locally integrable functions  $f$  such that

$$\|f\|_{BMO} = \sup_{x \in X, r > 0} \mu(B(x, r))^{-1} \int_{B(x, r)} |f(y) - f_{B(x, r)}| d\mu(y) < \infty$$

or

$$\|f\|_{BMO} = \inf_C \sup_{x \in X, r > 0} \mu(B(x, r))^{-1} \int_{B(x, r)} |f(y) - C| d\mu(y) < \infty.$$

**Definition 1.10.** Given a measurable function  $b$  the maximal commutator is defined by

$$M_b(f)(x) = \sup_{r > 0} \mu(B(x, r))^{-1} \int_{B(x, r)} |b(x) - b(y)| |f(y)| d\mu(y)$$

for all  $x \in X$ .

This operator plays an important role in the study of commutators of singular integral operators with BMO symbols.

**Definition 1.11.** Given a measurable function  $b$  the commutator of the Hardy-Littlewood maximal operator  $M$  and  $b$  is defined by

$$[M, b]f(x) = M(bf)(x) - b(x)Mf(x)$$

for all  $x \in X$ .

This operator arises, for example, when one tries to give a meaning to the product of a function in  $H^1$  and a function in BMO (which may not be a locally integrable function). Operators  $M_b$  and  $[M, b]$  essentially differ from each other. For example,  $M_b$  is a positive and sublinear operator, but  $[M, b]$  is neither positive nor sublinear. However, if  $b$  satisfies some additional conditions, then operator  $M_b$  controls  $[M, b]$ . The maximal operators and their commutators were widely investigated in various Morrey-type spaces (see, for example, [2, 5, 6, 13, 36, 40, 41]).

**Definition 1.12.** We define the  $BMO_{p, \omega}(X)$  ( $1 \leq p < \infty$ ) space as the set of all locally integrable functions  $f$  such that

$$\|f\|_{BMO_{p, \omega}} = \sup_{x \in X, r > 0} \frac{\|(f(\cdot) - f_{B(x, r)})\chi_{B(x, r)}\|_{L_{p, \omega}(X)}}{\|\omega\|_{L_p(B(x, r))}}$$

or

$$\|f\|_{BMO_{p, \omega}} = \sup_{x \in X, r > 0} \frac{1}{\mu(B(x, r))} \|(f(\cdot) - f_{B(x, r)})\chi_{B(x, r)}\|_{L_{p, \omega}(X)} \|\omega^{-1}\|_{L_{p'}(B(x, r))} < \infty.$$

The proof of the following theorem can be obtained similarly with the Corollary 4.5 in [24].

**Theorem 1.13.** Let  $1 \leq p < \infty$  and  $\omega$  be a Lebesgue measurable function. If  $\omega \in A_p(X)$ , then the norms  $\|\cdot\|_{BMO_{p, \omega}}$  and  $\|\cdot\|_{BMO}$  are mutually equivalent.

We will need the following lemma while proving our main theorems.

**Lemma 1.14.** [23] Let  $b \in BMO(X)$ . Then there is a constant  $C > 0$  such that

$$|b_{B(x, r)} - b_{B(x, t)}| \leq C \|b\|_{BMO} \ln \frac{t}{r} \quad \text{for } 0 < 2r < t,$$

where  $C$  is independent of  $b, x, r$ , and  $t$ .

Let  $L_{\infty,\omega}(\mathbb{R}_+)$  be the weighted  $L_\infty$ -space with the norm

$$\|g\|_{L_{\infty,\omega}(\mathbb{R}_+)} = \operatorname{ess\,sup}_{t>0} \omega(t)g(t).$$

We denote

$$\mathbb{A} = \left\{ \varphi \in \mathfrak{M}^+(\mathbb{R}_+; \uparrow) : \lim_{t \rightarrow 0+} \varphi(t) = 0 \right\}.$$

Let  $u$  be a continuous and non-negative function on  $\mathbb{R}_+$ . We define the supremal operator  $\bar{S}_u$  by

$$(\bar{S}_u g)(t) := \|u g\|_{L_\infty(0,t)}, \quad t \in (0, \infty).$$

The following theorem was proved in [9].

**Theorem 1.15.** [9] Suppose that  $v_1$  and  $v_2$  are nonnegative measurable functions such that  $0 < \|v_1\|_{L_\infty(0,t)} < \infty$  for every  $t > 0$ . Let  $u$  be a continuous nonnegative function on  $\mathbb{R}$ . Then the operator  $\bar{S}_u$  is bounded from  $L_{\infty,v_1}(\mathbb{R}_+)$  to  $L_{\infty,v_2}(\mathbb{R}_+)$  on the cone  $\mathbb{A}$  if and only if

$$\left\| v_2 \bar{S}_u \left( \|v_1\|_{L_\infty(0,\cdot)}^{-1} \right) \right\|_{L_\infty(\mathbb{R}_+)} < \infty.$$

We will use the following statement on the boundedness of the weighted Hardy operator

$$H_w g(t) := \int_0^t g(s)w(s)ds, \quad H_w^* g(t) := \int_t^\infty g(s)w(s)ds, \quad 0 < t < \infty,$$

where  $w$  is a weight.

The following theorem was proved in [19].

**Theorem 1.16.** [19] Let  $v_1, v_2$  and  $w$  be weights on  $(0, \infty)$  and  $v_1(t)$  be bounded outside a neighborhood of the origin. The inequality

$$\sup_{t>0} v_2(t) H_w^* g(t) \leq C \sup_{t>0} v_1(t) g(t)$$

holds for some  $C > 0$  for all non-negative and non-decreasing  $g$  on  $(0, \infty)$  if and only if

$$B := \sup_{t>0} v_2(t) \int_t^\infty \frac{w(s)ds}{\operatorname{ess\,sup}_{s<\tau<\infty} v_1(\tau)} < \infty.$$

**Theorem 1.17.** [19, 20] Let  $v_1, v_2$  and  $w$  be weights on  $(0, \infty)$  and  $v_1(t)$  be bounded outside a neighborhood of the origin. The inequality

$$\sup_{t>0} v_2(t) H_w g(t) \leq C \sup_{t>0} v_1(t) g(t) \tag{1}$$

holds for some  $C > 0$  for all non-negative and non-decreasing  $g$  on  $(0, \infty)$  if and only if

$$B := \sup_{t>0} v_2(t) \int_0^t \frac{w(s)ds}{\sup_{0<\tau<s} v_1(\tau)} < \infty.$$

Moreover, the value  $C = B$  is the best constant for (1).

## 2 Two-weighted inequalities for maximal commutators in generalized weighted Morrey spaces on spaces of homogeneous type

In this section we prove two-weighted inequalities for maximal commutators in generalized weighted Morrey spaces on spaces of homogeneous type.

**Lemma 2.1.** [1] Let  $b$  be any non-negative locally integrable function. Then

$$|[M, b]f(x)| \leq M_b(f)(x), \quad x \in X$$

holds for all  $f \in L_1^{loc}(X)$ .

**Theorem 2.2.** [1] Let  $b \in BMO(X)$ . Suppose that  $Y$  is a Banach space of measurable functions defined on  $X$ . Assume that  $M$  is bounded on  $Y$ . Then the operator  $M_b$  is bounded on  $Y$ , and the inequality

$$\|M_b f\|_Y \leq C \|b\|_{BMO} \|f\|_Y$$

holds with constant  $C$  independent of  $f$ .

**Corollary 2.3.** Let  $1 \leq p < \infty$ ,  $b \in BMO(X)$  and  $\omega \in A_p(X)$ , then the operator  $M_b$  is bounded in  $L_{p,\omega}(X)$ .

**Theorem 2.4.** Let  $1 < p < \infty$ ,  $0 < \delta < 1$ ,  $b \in BMO(X)$  and  $(\omega_1, \omega_2) \in \tilde{A}_p(X)$ ,  $\omega_1 \in A_p(X)$ , then the operator  $M_b$  is bounded from  $L_{p,\omega_1^\delta}(X)$  to  $L_{p,\omega_2^\delta}(X)$ .

*Proof.* Let  $f \in L_{p,\omega_1^\delta}(X)$ ,  $b \in BMO(X)$ . The inequality ([1], Corollary 1.11), is valid

$$M_b f(x) \leq C \|b\|_{BMO} M^2 f(x).$$

By the this inequality, Corollary 1.7, Corollary 2.3 and  $(\omega_1, \omega_2) \in \tilde{A}_p(X)$ ,  $\omega_1 \in A_p(X)$  conditions, we have

$$\begin{aligned} \|M_b f\|_{L_{p,\omega_2^\delta}(X)} &\leq C \|b\|_{BMO} \|M^2 f\|_{L_{p,\omega_2^\delta}(X)} \\ &\leq C \|b\|_{BMO} \|M f\|_{L_{p,\omega_1^\delta}(X)} \leq C_1 \|b\|_{BMO} \|f\|_{L_{p,\omega_1^\delta}(X)}, \end{aligned}$$

where  $M^2 f(x) = M(M f(x))$ .

**Theorem 2.5.** Let  $1 < p < \infty$ ,  $0 < \delta < 1$ ,  $b \in BMO(X)$  and  $(\omega_1, \omega_2) \in \tilde{A}_p(X)$ ,  $\omega_1, \omega_2 \in A_p(X)$ , then

$$\|M_b f\|_{L_{p,\omega_2^\delta}(B(x,r))} \leq C \|b\|_{BMO} \|\omega_2^\delta\|_{L_p(B(x,r))} \sup_{t>r} \left(1 + \ln \frac{t}{r}\right) \frac{\|f\|_{L_{p,\omega_1^\delta}(B(x,t))}}{\|\omega_2^\delta\|_{L_p(B(x,t))}}, \quad (1)$$

for every  $f \in L_{p,\omega_1^\delta}(X)$ , where  $C$  does not depend on  $f, x$  and  $r$ .

*Proof.* We represent function  $f$  as

$$f = f_1 + f_2, \quad f_1(y) = f(y) \chi_{B(x,2kr)}(y), \quad f_2(y) = f(y) \chi_{X \setminus B(x,2kr)}(y), \quad r > 0,$$

where  $k$  is the constant from the quasi-triangle inequality and have

$$\|M_b f\|_{L_{p,\omega_2^\delta}(B(x,r))} \leq \|M_b f_1\|_{L_{p,\omega_2^\delta}(B(x,r))} + \|M_b f_2\|_{L_{p,\omega_2^\delta}(B(x,r))}.$$

By Theorem 2.4 we obtain

$$\begin{aligned} \|M_b f_1\|_{L_{p,\omega_2^\delta}(B(x,r))} &\leq \|M_b f_1\|_{L_{p,\omega_2^\delta}(X)} \\ &\leq C \|b\|_{BMO} \|f_1\|_{L_{p,\omega_1^\delta}(X)} = C \|b\|_{BMO} \|f\|_{L_{p,\omega_1^\delta}(B(x,2kr))}, \end{aligned} \quad (2)$$

where  $C$  does not depend on  $f$ . From (2) we get

$$\|M_b f_1\|_{L_{p,\omega_2^\delta}(B(x,r))} \leq C \|b\|_{BMO} \|\omega_2^\delta\|_{L_p(B(x,r))} \sup_{t>r} \left(1 + \ln \frac{t}{r}\right) \frac{\|f\|_{L_{p,\omega_1^\delta}(B(x,t))}}{\|\omega_2^\delta\|_{L_p(B(x,t))}}, \quad (3)$$

which is easily obtained from the fact that  $\|f\|_{L_{p,\omega_1^\delta}(B(x,2kr))}$  is non-decreasing in  $r$ , therefore  $\|f\|_{L_{p,\omega_1^\delta}(B(x,2kr))}$  on the right-hand side of (2) is dominated by the right-hand side of (3).

For  $y \in B(x,r)$  we get

$$\begin{aligned} M_b f_2(y) &= \sup_{t>0} \mu(B(y,t))^{-1} \int_{B(y,t)} |b(y) - b(z)| |f_2(z)| d\mu(z) \\ &= \sup_{t>0} \mu(B(y,t))^{-1} \int_{B(x,2kr) \cap B(y,t)} |b(y) - b(z)| |f(z)| d\mu(z) \\ &\leq C \sup_{t>2kr} \mu(B(y,t))^{-1} \int_{B(x,t)} |b(y) - b(z)| |f(z)| d\mu(z) \\ &\leq \sup_{t>r} \mu(B(y,t))^{-1} \int_{B(x,t)} |b(z) - b_{B(x,t)}| |f(z)| d\mu(z) \\ &\quad + \sup_{t>r} \mu(B(y,t))^{-1} \int_{B(x,t)} |b(y) - b_{B(x,t)}| |f(z)| d\mu(z) = I_1 + I_2. \end{aligned}$$

By Hölder inequality and Theorem 1.13 we obtain

$$\begin{aligned} I_1 &= \sup_{t>r} \mu(B(y,t))^{-1} \int_{B(x,t)} |b(z) - b_{B(x,t)}| |f(z)| d\mu(z) \\ &\leq C \sup_{t>r} \mu(B(y,t))^{-1} \|f\|_{L_{p,\omega_1^\delta}(B(x,t))} \|b(\cdot) - b_{B(x,t)}\|_{L_{p',\omega_1^{-\delta}}(B(x,t))} \\ &\leq C \|b\|_{BMO} \sup_{t>r} t^{-N} \|f\|_{L_{p,\omega_1^\delta}(B(x,t))} \|\omega_1^{-\delta}\|_{L_{p'}(B(x,t))}. \end{aligned}$$

To estimate  $I_2$ , by Lemma 1.14 we get

$$\begin{aligned} I_2 &= \sup_{t>r} \mu(B(y,t))^{-1} |b(y) - b_{B(x,t)}| \int_{B(x,t)} |f(z)| d\mu(z) \\ &\leq C M_b \chi_{B(x,r)}(y) \sup_{t>r} t^{-N} \|f\|_{L_{p,\omega_1^\delta}(B(x,t))} \|\omega_1^{-\delta}\|_{L_{p'}(B(x,t))} \\ &\quad + C \|b\|_{BMO} \sup_{t>r} t^{-N} \ln \frac{t}{r} \|f\|_{L_{p,\omega_1^\delta}(B(x,t))} \|\omega_1^{-\delta}\|_{L_{p'}(B(x,t))}. \end{aligned}$$



From Theorem 2.2 we have

$$\begin{aligned}
 \|M_b f_2\|_{L_{p,\omega_2^\delta}(B(x,r))} &\leq \|I_1\|_{L_{p,\omega_2^\delta}(B(x,r))} + \|I_2\|_{L_{p,\omega_2^\delta}(B(x,r))} \\
 &\leq C\|b\|_{BMO}\|\omega_2^\delta\|_{L_p(B(x,r))} \sup_{t>r} \left(1 + \ln \frac{t}{r}\right) \frac{\|f\|_{L_{p,\omega_1^\delta}(B(x,t))}}{\|\omega_2^\delta\|_{L_p(B(x,t))}} \\
 &\quad + C\|M_b \chi_{B(x,r)}\|_{L_{p,\omega_2^\delta}(B(x,r))} \sup_{t>r} \left(1 + \ln \frac{t}{r}\right) \frac{\|f\|_{L_{p,\omega_1^\delta}(B(x,t))}}{\|\omega_2^\delta\|_{L_p(B(x,t))}} \\
 &\leq C\|b\|_{BMO}\|\omega_2^\delta\|_{L_p(B(x,r))} \sup_{t>r} \left(1 + \ln \frac{t}{r}\right) \frac{\|f\|_{L_{p,\omega_1^\delta}(B(x,t))}}{\|\omega_2^\delta\|_{L_p(B(x,t))}}. \tag{4}
 \end{aligned}$$

Then from (3) and (4) we obtain (1).

**Theorem 2.6.** Let  $1 < p < \infty$ ,  $0 < \delta < 1$ ,  $b \in BMO(X)$ ,  $(\omega_1, \omega_2) \in \tilde{A}_p(X)$ ,  $\omega_1, \omega_2 \in A_p(X)$  and the functions  $\varphi_1(x, r)$ ,  $\varphi_2(x, r)$  satisfy the condition

$$\sup_{t>r} \left(1 + \ln \frac{t}{r}\right) \frac{\operatorname{ess\,inf}_{t<s<\infty} \varphi_1(x, s) \|\omega_1^\delta\|_{L_p(B(x,s))}}{\|\omega_2^\delta\|_{L_p(B(x,t))}} \leq C\varphi_2(x, r), \tag{5}$$

where  $C$  does not depend on  $x$  and  $t$ .

Then the operator  $M_b$  is bounded from the space  $\mathcal{M}_{\omega_1^\delta}^{p,\varphi_1}(X)$  to the space  $\mathcal{M}_{\omega_2^\delta}^{p,\varphi_2}(X)$ .

*Proof.* Let  $f \in \mathcal{M}_{\omega_1^\delta}^{p,\varphi_1}(X)$ . Let us take  $v_2 = \frac{1}{\varphi_2(x,t)}$ ,  $g = \|f\|_{L_{p,\omega_1^\delta}(B(x,t))}$ ,  $u = (1 + \ln \frac{t}{r}) \|\omega_2^\delta\|_{L_p(B(x,t))}^{-1}$ ,  $v_1 = \frac{1}{\varphi_1(x,t) \|\omega_1^\delta\|_{L_p(B(x,t))}}$  in Theorem 1.16. Also by using Theorem 2.5 and the inequality (5) we get

$$\begin{aligned}
 &\|M_b f\|_{\mathcal{M}_{\omega_2^\delta}^{p,\varphi_2}(X)} \\
 &\leq C\|b\|_{BMO} \sup_{x \in X, r>0} \frac{\|\omega_2^\delta\|_{L_p(B(x,r))}}{\varphi_2(x, r) \|\omega_2^\delta\|_{L_p(B(x,r))}} \sup_{t>r} \left(1 + \ln \frac{t}{r}\right) \frac{\|f\|_{L_{p,\omega_1^\delta}(B(x,t))}}{\|\omega_2^\delta\|_{L_p(B(x,t))}} \\
 &\leq C\|b\|_{BMO} \sup_{x \in X, r>0} \frac{1}{\varphi_1(x, r) \|\omega_1^\delta\|_{L_p(B(x,r))}} \|f\|_{L_{p,\omega_1^\delta}(B(x,r))} = C\|b\|_{BMO} \|f\|_{\mathcal{M}_{\omega_1^\delta}^{p,\varphi_1}(X)},
 \end{aligned}$$

which completes the proof.

### 3 Declarations

#### Funding

The research of Javanshir J. Hasanov was supported by The Turkish Scientific and Technological Research Council (TUBITAK, programme 2221).

#### Competing Interests

The authors declare that they have no competing interests.

#### Ethical Approval

Not applicable.

## Authors' Contributions

All authors contributed equally. All the authors read and approved the final manuscript.

## Availability Data and Materials

Not applicable.

## Acknowledgements

The authors would like to thank the anonymous reviewer for his/her comments.

## References

- [1] M. Ağcayazı, A. Gogatishvili, K. Koca and R. Mustafayev, *A note on maximal commutators and commutators of maximal functions*, J. Math. Soc. Japan. **67** (2015), no. 2, 581-593.
- [2] M. Ağcayazı, A. Gogatishvili and R. Mustafayev, *Weak-type estimates in Morrey spaces for maximal commutator and commutator of maximal function*, Tokyo J. Math. **41** (2018), no. 1, 193-218.
- [3] C. Avşar, C. Aykol, J.J. Hasanov and A.M. Musayev, *Two-weight inequalities for Riesz potential and its commutators on weighted global Morrey-type spaces  $\mathcal{GM}_{\omega}^{p,\theta,\varphi}(\mathbb{R}^n)$* , Advanced Studies: Euro-Tbilisi Math. J. **16** (2023), no. 1, 33-50.
- [4] A. Aydoğdu and C. Aykol, *Two-weighted inequalities for maximal operator in generalized weighted Morrey spaces on spaces of homogeneous type*, Baku Math. J. **2** (2023), no. 1, 113-121.
- [5] C. Aykol, H. Armutçu and M.N. Omarova, *Maximal commutator and commutator of maximal function on modified Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **36** (2016), no. 1, 29-35.
- [6] C. Aykol, X.A. Badalov and J.J. Hasanov, *Maximal and singular operators in the local "complementary" generalized variable exponent Morrey spaces on unbounded sets*, Quaest. Math. **43** (2020), no. 10, 1487-1512.
- [7] C. Aykol, J.J. Hasanov and Z.V. Safarov, *Two weighted inequalities for Riesz potential and its commutator in generalized weighted Morrey spaces*, Mat. Vesnik **75** (2023), no. 1, 37-49.
- [8] C. Aykol, J.J. Hasanov and Z.V. Safarov, *A characterization of two-weighted inequalities for maximal, singular operators and their commutators in generalized weighted Morrey spaces*, Funct. Approx. Comment. Math. **67** (2022), no. 2, 145-167.
- [9] V. Burenkov, A. Gogatishvili, V.S. Guliyev and R. Mustafayev, *Boundedness of the fractional maximal operator in local Morrey-type spaces*, Complex Var. Elliptic Equ. **55** (2010), no. 8-10, 739-758.
- [10] R.R. Coifman and G. Weiss, *Analyse harmonique non-commutative sur certain espaces homogenes*, in "Lecture Notes in Math.," No. 242, Springer-Verlag, Berlin, (1971).
- [11] D. Cruz-Urbe, *New proofs of Two-weight norm inequalities for the maximal operator*, Georgian Math. J. (2000), no. 7, 33-42.
- [12] D. Cruz-Urbe, *Two weight norm inequalities for fractional integral operators and commutators*, Advanced Courses of Mathematical Analysis VI, World Sci. Publ., Hackensack, NJ, (2017), 25-85.
- [13] G. Di Fazio and M. A. Ragusa, *Commutators and Morrey spaces*, Bollettino U.M.I. **7** 5-A (1991), 323-332.
- [14] D.E. Edmunds and V.M. Kokilashvili, *Two weighted inequalities for singular integrals*, Canad. Math. Bull. **38** (1995), no. 3, 295-303.
- [15] V.S. Guliyev, *Integral operators on function spaces on the homogeneous groups and on domains in  $\mathbb{G}$* , (Russian) Doctor's degree dissertation, Moscow, Mat. Inst. Steklov, 1-329 (1994).
- [16] V.S. Guliyev, *Function spaces, integral operators and two weighted inequalities on homogeneous groups*, Some applications. (Russian) Baku, 1-332 (1999).
- [17] V.S. Guliyev, *Boundedness of the maximal, potential and singular operators in the generalized Morrey spaces*, J. Inequal. Appl. Art. **1** (2009).
- [18] V.S. Guliyev, *Generalized weighted Morrey spaces and higher order commutators of sublinear operators*, Eurasian Math. J. **3** (2012), no. 3, 33-61.
- [19] V.S. Guliyev, *Generalized local Morrey spaces and fractional integral operators with rough kernel*, J. Math. Sci. (N. Y.) **193** (2013), no. 2, 211-227.
- [20] V.S. Guliyev, *Local generalized Morrey spaces and singular integrals with rough kernel*, Azerb. J. Math. **3** (2013), no. 2, 79-94.
- [21] V.S. Guliyev, T. Karaman, R.Ch. Mustafayev and A. Serbetci, *Commutators of sublinear operators generated by Calderón-Zygmund operator on generalized weighted Morrey spaces*, Czechoslovak Math. J. **60** (2014), no. 1, 365-386.

- [22] D.D. Haroske and L. Skrzypczak, *Embeddings of weighted Morrey spaces*, Math. Nachr. **290** (2017), no. 7, 1066-1086.
- [23] T. Heikkinen, J. Kinnunen, J. Nuutinen and H. Tuominen, *Mapping properties of the discrete fractional maximal operator in metric measure spaces*, Kyoto J. Math. **53** (2013), no. 3, 693–712.
- [24] K.P. Ho, *Singular integral operators, John-Nirenberg inequalities and Tribel-Lizorkin type spaces on weighted Lebesgue spaces with variable exponents*, Revista De La Union Matematica Argentina **57** (2016), no. 1, 85-101.
- [25] T. Karaman, V.S. Guliyev and A. Serbetci, *Boundedness of sublinear operators generated by Calderón-Zygmund operators on generalized weighted Morrey spaces*, An. Stiint. Univ. Al. I. Cuza Iasi. Mat. (N. S.) **60** (2014), no. 1, 227-244.
- [26] V. Kokilashvili and A. Meskhi, *Two-weight inequalities for fractional maximal functions and singular integrals in  $L^{p(\cdot)}$  spaces*, J. Math. Sci. (N.Y.) **173** (2011), no. 6, 656-673.
- [27] Y. Komori and S. Shirai, *Weighted Morrey spaces and a singular integral operator*, Math. Nachr. **282** (2009), no. 2, 219-231.
- [28] M. T. Lacey, E. T. Sawyer, and I. Uriarte-Tuero, *A characterization of two weight norm inequalities for maximal singular integrals with one doubling measure*, Anal. PDE **5** (2012), no. 1, 1-60.
- [29] T. Mizuhara, *Boundedness of some classical operators on generalized Morrey spaces*, Harmonic Analysis (S. Igari, Editor), ICM 90 Satellite Proceedings, Springer - Verlag, Tokyo 183-189 (1991).
- [30] C.B. Morrey, *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc. **43** (1938), 126-166.
- [31] B. Muckenhoupt, *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc. **165**(1972), 207-226.
- [32] E. Nakai, *Hardy-Littlewood maximal operator, singular integral operators and Riesz potentials on generalized Morrey spaces*, Math. Nachr. **166** (1994), 95-103.
- [33] S. Nakamura, Y. Sawano, and H. Tanaka, *The fractional operators on weighted Morrey spaces*, J. Geom. Anal. **28** (2018), no. 2, 1502-1524.
- [34] C.J. Neugebauer, *Inserting  $A_p$ -weights*, Proc. Amer. Math. Soc. **87** (1983), no. 4, 644-648.
- [35] J. Pan and W. Sun, *Two weight norm inequalities for fractional maximal functions and fractional integral operators on weighted Morrey spaces*, Math. Nachr. **293** (2020), no. 5, 970-982.
- [36] M.A. Ragusa, *Regularity of solutions of divergence form elliptic equations*, Proc. Amer. Math. Soc. **128** (2000), no. 1, 533-540.
- [37] N. Samko, *Weighted Hardy and singular operators in Morrey spaces*, J. Math. Anal. Appl. **350** (2009), no. 1, 56-72.
- [38] Y. Sawano, *A thought on generalized Morrey spaces*, J. Indonesian Math. Soc. **25** (2019), no. 3, 210-281 .
- [39] H. Tanaka, *Two-weight norm inequalities on Morrey spaces*, Ann. Acad. Sci. Fenn. Math. **40** (2015), no. 2, 773-791.
- [40] L.W. Wang, *The commutators of multilinear maximal and fractional-type operators on central Morrey spaces with variable exponent*, J. Funct. Spaces **2022**, art.n. 4875460, (2022).
- [41] T.L. Yee, K.L. Cheung , K.P. Ho, *Integral operators on local Orlicz-Morrey spaces*, Filomat **36** (2022), no. 4, 1231-1243.